



MONASH
University



COEPP

ARC Centre of Excellence for
Particle Physics at the Terascale

MoCA
Monash Centre for Astrophysics



GAMBIT

update

Csaba Balázs

for the GAMBIT Community

2021 Oct 15 Taipei



outline

global fits

GAMBIT

community

main features

code structure

version 2.0

the future



global fitting

Why?

standard models emerge from a set of competing theories


the simplest theory best fitting the most data becomes standard model

global fitting is to quantify “simplest”, “best”, and “most” above



global fitting

What?



global fitting

How?

main steps

- establish model hierarchy
- Lagrangian → spectrum
- spectrum → observables
- parameter space sampling
- statistical treatment

main challenges

- many models (& parameters)
- spectrum auto-generation
- fast backends, auto-generation
- efficiency: need for speed
- rigorous, meaningful inference

these steps and challenges can be tackled “model independently”



open-source **global** fitting framework

modular and flexible architecture

models **beyond** the standard

sophisticated statistical **inference**

plug&play **tools** to calc observables



Global And Modular BSM Inference Tool

gambit.hepforge.org

EPJC 77 (2017) 784, arXiv:1705.07908

- open-source code to calculate observables and likelihoods for generic Beyond the Standard Model(s) theories
- designed to allow easy definition of new models, observables, likelihoods, samplers and backend physics codes
- extensive
 - model database
 - observable calculators
 - data libraries
- various sampling and stat options
- fast likelihood calculators (LHC...)
- massively parallel



Community

subset

70+ community members


F Agocs, V Ananyev, P Athron, C Balázs, A Beniwal, J Bhom, S Bloor, T Bringmann, A Buckley, J-E Camargo-Molina, C Chang, M Chrzaszcz, J Conrad, J Cornell, M Danner, J Edsjö, B Farmer, A Fowlie, T Gonzalo, P Grace, W Handley, J Harz, S Hoof, S Hotinli, F Kahlhoefer, N Avis Kozar, A Kvellestad, P Jackson, A Ladhu, N Mahmoudi, G Martinez, MT Prim, F Rajec, A Raklev, J Renk, C Rogan, R Ruiz, I Sáez Casares, N Serra, A Scaffidi, P Scott, P Stöcker, W Su, J Van den Abeele, A Vincent, C Weniger, M White, Y Zhang ...

from 14+ countries, 29+ institutes



Community

subset



Community

linkages

members of many experiments

ATLAS, Belle-II, CLiC, CMS, CTA, Fermi-LAT, DARWIN, IceCube, LHCb,
SHiP, XENON

authors of numerous theory codes

BubbleProfiler, Capt'n General, DarkSUSY, DDCalc, Diver, FlexibleSUSY,
gamlike, GM2Calc, HEPLike, IsaJet, nulike, PhaseTracer, PolyChord,
Rivet, SOFTSUSY, SuperIso, SUSY-AI, xsec, Vevacious, WIMPSim...



GAMBIT features

global and modular

- diverse BSM model database SM+SS, EFTs, 2HDMs, MSSM63, axions, RHNs, cosmo...
- changeable model assumptions for cosmology, astro-, particle-, nuclear physics...
- composite likelihood consistent combination of searches, uncertainties, nuisances...
- built-in experimental likelihoods LEP, ATLAS, CMS, LHCb, DM searches...
- sampling algorithms (ensemble) MCMC, T-walk, diff. evolution, particle swarm, nested...
- auto dependency resolution ID functions, optimize execution order! before run
- dual-level parallel execution mixed-mode MPI+openMP, mostly auto, scale 10k+ cores
- many interfaced backends observable calculators for cosmology, astrophysics, collider, precision, flavor... (list on later slide)
- diskless generalization of various Les Houches Accords




GAMBIT features

flexible and extendable

- fast definition of new models, data sets, sampling methods
- plug&play theory tools auto-download, configure, compile, dynamically link
- easily switch between backends calculating the same quantities
- C/C++, Python, Fortran, Mathematica interfaces for backends
- BOSS dynamic loading of C++ classes from backend shared libraries!
- all-in or module standalone modes easily implemented from single cmake script
- YAML input model, parameters, observables, sampler, stat. inference
- customizable output streams ASCII, HDF5, databases...
- advanced statistical inference parameter estimation, Bayesian model comparison
- available as docker plugin <https://gambit.hepforge.org/source>





Models

hierarchical model database


- models defined by their parameters and relations to each other
- models can inherit from (be subspaces of) parent models
- child models can be automatically translated to ancestor models
- database examples:
 - standard model
 - nuclear uncertainties
 - two-Higgs doublet models
 - Higgs portal dark matter models
 - dark matter halo models
 - right-handed neutrino models
 - effective field theory dark matter models...



Models

model database examples


- axions and ALPs



Models

model database examples

- MSSM



Modules model independent, interdependent physics structures

- ColliderBit: event gen., fast sim., Z, H obs.s, search limits... arXiv:1705.07919
- DarkBit: DM abundance, direct-, indirect detection... arXiv:1705.07920
- DecayBit: SM & NP (SUSY...) decay widths, BRs... arXiv:1705.07936
- FlavBit: NP (SUSY...) 100s of flavor obs.s, rare decays... arXiv:1705.07933
- PrecisionBit: EW precision observables, g-2... arXiv:1705.07936
- SpecBit: SM & NP masses, mixings, couplings, RGEs... arXiv:1705.07936
- ScannerBit: sampling, parameter est., model comparison... arXiv:1705.07959
- NeutrinoBit: neutrino observables, likelihoods, RHNs... arXiv:1908.02302
- CosmoBit: Λ CDM+, inflation, neutrinos, axions... arXiv:2009.03286
- GUM: auto-generation (spectrum, interfaces, observables)... arXiv:2107.00030



Backends

observable calculators for cosmology,
astrophysics, collider, precision, flavor...


- AlterBBN
- CalcHEP
- Capt'n General
- CLASS
- DarkAges
- DDCalc
- DarkSUSY
- FeynHiggs
- FeynRules
- Flavio
- FlexibleSUSY
- gamLike
- GM2Calc
- HepLike
- HiggsBounds
- HiggsSignals
- MadGraph
- micrOMEGAs
- MontePython
- MultiModeCode
- nulike
- pic
- Pythia
- SARAH
- Spheno
- SUSYHD
- SUSY-HIT
- SuperIso
- Vevacious
- ...



GAMBIT


run overview

- user chooses model, observables, sampler
- using graph-theory GAMBIT constructs a dependency tree to optimize the calculation
- GAMBIT samples para. space by calling the necessary module and backend functions for each parameter point



Dependency resolution

for CMSSM



Dependency resolution

for CMSSM

model parameter translations

precision calculations


LEP rates+likelihoods

decays

LHC observables and likelihoods

DM abundance, direct, indirect searches

flavour
physics



Derivation

dependencies constructed
dynamically at run-time using
graph-theoretic methods to solve
for required observables,
backends, evaluation order, etc.


precision calculations

flavor
physics


LEP

Log-likelihoods

indirect, direct, and
indirect searches



S



GAMBIT 2.0


GAMBIT Universal Model

from Lagrangian to likelihood




- auto-generating GAMBIT code and interfaces to backends
- from Lagrangian-level input use FeynRules, SARAH, MadGraph, CalcHEP to generate GAMBIT model, collider, dark matter, decay and spectrum code
- GUM also writes C++ GAMBIT interfaces to SPheno, micrOMEGAs, Pythia, Vevacious
- arXiv:2107.00030 worked example: addition of a Majorana fermion simplified dark matter model with a scalar mediator to GAMBIT via GUM, and carry out a corresponding fit

















GAMBIT 2

GUM schematics





some results



Dark matter effective field theory arXiv:2106.02056

- general DMEFT: all DM-SM effective interactions up to dimension n
- in principle, a global fit of the general theory is possible with GAMBIT
- we focus on the 16D space of Dirac DM interactions with SM quarks and gluons:

$$\mathcal{L}_\chi = \bar{\chi} \left(i\cancel{D} - \cancel{m}_\chi \right) \chi + \sum_{a,d} \frac{\mathcal{C}_a^{(d)}}{\Lambda^{d-4}} \mathcal{Q}_a^{(d)}$$



Dark matter effective field theory arXiv:2106.02056

$$\mathcal{L}_\chi = \bar{\chi} (i\not{\partial} - m_\chi) \chi + \sum_{a,d} \frac{\mathcal{C}_a^{(d)}}{\Lambda^{d-4}} Q_a^{(d)}$$

Direct detection signals:

Spin-independent – not suppressed

Spin-independent – suppressed

Spin-dependent – not suppressed

Spin-dependent – suppressed

$$Q_{1,q}^{(6)} = (\bar{\chi} \gamma_\mu \chi)(\bar{q} \gamma^\mu q)$$

$$Q_1^{(7)} = \frac{\alpha_s}{12\pi} (\bar{\chi} \chi) G^{a\mu\nu} G^a_{\mu\nu}$$

$$Q_{6,q}^{(7)} = m_q (\bar{\chi} i \gamma_5 \chi)(\bar{q} q)$$

$$Q_{2,q}^{(6)} = (\bar{\chi} \gamma_\mu \gamma_5 \chi)(\bar{q} \gamma^\mu q)$$

$$Q_2^{(7)} = \frac{\alpha_s}{12\pi} (\bar{\chi} i \gamma_5 \chi) G^{a\mu\nu} G^a_{\mu\nu}$$

$$Q_{7,q}^{(7)} = m_q (\bar{\chi} \chi)(\bar{q} i \gamma_5 q)$$

$$Q_{3,q}^{(6)} = (\bar{\chi} \gamma_\mu \chi)(\bar{q} \gamma^\mu \gamma_5 q)$$

$$Q_3^{(7)} = \frac{\alpha_s}{8\pi} (\bar{\chi} \chi) G^{a\mu\nu} \tilde{G}^a_{\mu\nu}$$

$$Q_{8,q}^{(7)} = m_q (\bar{\chi} i \gamma_5 \chi)(\bar{q} i \gamma_5 q)$$

$$Q_{4,q}^{(6)} = (\bar{\chi} \gamma_\mu \gamma_5 \chi)(\bar{q} \gamma^\mu \gamma_5 q)$$

$$Q_4^{(7)} = \frac{\alpha_s}{8\pi} (\bar{\chi} i \gamma_5 \chi) G^{a\mu\nu} \tilde{G}^a_{\mu\nu}$$

$$Q_{9,q}^{(7)} = m_q (\bar{\chi} \sigma^{\mu\nu} \chi)(\bar{q} \sigma_{\mu\nu} q)$$

$$Q_{5,q}^{(7)} = m_q (\bar{\chi} \chi)(\bar{q} q)$$

$$Q_{10,q}^{(7)} = m_q (\bar{\chi} i \sigma^{\mu\nu} \gamma_5 \chi)(\bar{q} \sigma_{\mu\nu} q)$$



- the scale of new physics Λ is an independent parameter
- relic density calculation requires $\Lambda > 2m_\chi$
- if $\Lambda >$ scale probed by other experiments, we compute $\ln \mathcal{L}_{\text{experiment}}$
- otherwise, we set $\ln \mathcal{L}_{\text{experiment}} = 0$
- for LHC we smoothly cut off the spectrum to suppress events with *missing E_T* $> \Lambda$



Other key innovations

arXiv:2106.02056

- direct detection
 - DirectDM: fully automated RG evolution from Λ to low energies and matching to non-relativistic effective operators at hadronic scale
 - DDCalc: large database of direct detection constraints for arbitrary DM-nucleon interactions including astrophysical and nuclear uncertainties
- LHC constraints (ColliderBit)
 - monojet analyses: ATLAS 139/fb (full Run 2 dataset) + CMS 36/fb
 - fast profiling of LHC nuisance parameters



Other key innovations


arXiv:2106.02056

- indirect detection
 - DarkSUSY: highly accurate relic density calculation
 - GUM: automated calculation of cross sections and γ -ray spectra
 - CosmoBit: CMB constraints on energy injection from DM annihilation
 - Capt'n General: solar capture + neutrinos with arbitrary DM-nucleon interactions
- interface between all of these and Diver differential sampler fully automated in GAMBIT



DMEFT results


- new physics scale Λ :
- EFT valid for all constraints
- most experiments are insensitive
- constraints driven by relic density requirement
- Λ comparable to LHC energies
- strong LHC constraints
- Λ below LHC energies
- large viable parameter space



DMEFT results

- light DM ($m_\chi < 100$ GeV) is viable if both of the following hold:
 - χ has CP-violating interactions OR is very sub-dominant
 - Λ is so low that EFT breaks down entirely at LHC \Rightarrow LHC would (probably) be sensitive to mediator
- significant to model building, suggests light WIMPs require light mediator (e.g. dark photon)


light DM particles



DMEFT results

- demanding χ to be all of DM pushes viable parameter space to large m_χ
- leads to detectable signals at LZ
- mostly due to loop-induced operator mixing $Q(6)3,q \rightarrow Q(6)1,q$
- could be spoilt by including other effective operators (e.g. leptons, non-MFV) \Rightarrow interesting avenue for future investigation

heavy DM particles



public results available on
zenodo.cern.ch

- parameter point samples
- input files for all scans
- example plotting routines

links at gambit.hepforge.org/pubs

The screenshot shows the Zenodo search results page with the following details:

- Dataset 1:** **Supplementary Data: Impact of vacuum stability, perturbativity and XENON1T on global fits of Z2 and Z3 scalar singlet dark matter (arXiv:1806.11281)**
Uploaded on June 29, 2018 (v1) | Dataset | Open Access | View
- Dataset 2:** **Supplementary Data: Status of the scalar singlet dark matter model (arXiv:1705.07931)**
Uploaded on August 22, 2017 (v2) | Dataset | Open Access | View
- Dataset 3:** **Supplementary Data: A global fit of the MSSM with GAMBIT (arXiv:1705.07917)**
Uploaded on August 15, 2017 (v2) | Dataset | Open Access | View
- Dataset 4:** **Supplementary Data: Global fits of GUT-scale SUSY models with GAMBIT (arXiv:1705.07935)**
Uploaded on August 15, 2017 (v2) | Dataset | Open Access | View

The datasets are described as containing data for scalar singlet dark matter models, the MSSM, and GUT-scale SUSY models, with links to arXiv preprints and GAMBIT "Round 1" papers.



- more GAMBIT: more models, more observables, more automation, more data, more stats...
- machine learning of cross sections, cosmic ray fluxes...
- observable calculation on parallel GPUs
- ColliderBit Solo
- papers focusing on light SUSY, 2HDMs, DM direct detection, neutrinos, axions, leptoquarks, ...
- getting long-haired white cat, taking over the world



summary

- global fitting paves the way to the next standard model(s)
- GAMBIT is an open source, flexible, modular global fitting framework
- GUM significantly enhances the capabilities of GAMBIT 2.0
- GAMBIT version 2.1 just landed https://github.com/GambitBSM/gambit_2.1
- over a dozen physics papers during the last few years and many more to come
- stay tuned for much more...





backup slides

getting started

- clone git repo github.com/GambitBSM/gambit_2.0 or
- download tarballs <https://gambit.hepforge.org/source> or
- get pre-compiled version docker
- see quick start guides in arXiv:1705.07908 and arXiv:2107.00030



adding a new model to GAMBIT manually

1. Add the model to the **model hierarchy**:

- Choose a model name, and declare any **parent model**
- Declare the model's parameters
- Declare any **translation function** to the parent model

```
#define MODEL NUHM1
#define PARENT NUHM2
START_MODEL
DEFINEPARS(M0,M12,mH,A0,TanBeta,SignMu)
INTERPRET_AS_PARENT_FUNCTION(NUHM1_to_NUHM2)
#undef PARENT
#undef MODEL
```

2. Write the translation function as a standard C++ function:

```
void MODEL_NAMESPACE::NUHM1_to_NUHM2 (const ModelParameters &myP, ModelParameters &targetP)
{
    // Set M0, M12, A0, TanBeta and SignMu in the NUHM2 to the same values as in the NUHM1
    targetP.setValues(myP, false);
    // Set the values of mHu and mHd in the NUHM2 to the value of mH in the NUHM1
    targetP.setValue("mHu", myP["mH"]);
    targetP.setValue("mHd", myP["mH"]);
}
```

3. If needed, declare that existing module functions work with the new model, or add new functions that do.



adding a new observable/likelihood to GAMBIT manually

Adding a new module function is easy:

1. Declare the function to GAMBIT in a module's **rollcall header**

- Choose a capability
- Declare any **backend requirements**
- Declare any **dependencies**
- Declare any specific **allowed models**
- other more advanced declarations also available

```
#define MODULE FlavBit                                // A tasty GAMBIT module.  
START_MODULE  
  
#define CAPABILITY Rmu                               // Observable: BR(K->mu nu)/BR(pi->mu nu)  
START_CAPABILITY  
    #define FUNCTION SI_Rmu                         // Name of a function that can compute Rmu  
    START_FUNCTION(double)                        // Function computes a double precision result  
    BACKEND_REQ(Kmunu_pimunu, (my_tag), double, (const parameters*)) // Needs function from a backend  
    BACKEND_OPTION( (SuperIso, 3.6), (my_tag) )      // Backend must be SuperIso 3.6  
    DEPENDENCY(SuperIso_modelinfo, parameters)     // Needs another function to calculate SuperIso info  
    ALLOW_MODELS(MSSM63atQ, MSSM63atMGUT)          // Works with weak/GUT-scale MSSM and descendants  
    #undef FUNCTION  
#undef CAPABILITY
```

2. Write the function as a standard C++ function
(one argument: the result)



GAMBIT results related to dark matter

EFTs: SM + scalar, fermion, vector singlet (simplest DM model)

EPJ C77 (2017) 8 568 arXiv:1705.07931

EPJ C78 (2018) 10 830 arXiv:1806.11281

EPJ C79 (2019) 1 38 arXiv:1808.10465

arXiv:2106.02056

axions, axion-like particles (QCD axion, DFSZ, KSVZ, generic ALP)

JHEP 1903 (2019) 191 arXiv:1810.07192

arXiv:2006.03489

constrained SUSY: CMSSM, NUHM1, NUHM2 (GUT scale BCs)

EPJ C77 (2017) 12 824 arXiv:1705.07935

low-dim SUSY: MSSM-7, MSSM-EW (weak scale BCs)

EPJ C77 (2017) 12 879 arXiv:1705.07917

EPJ C79 (2019) 5 395 arXiv:1809.02097

more EFTs, more ALPs, more SUSY and other models in prep

